



FORAGE FIRST®

Top Tim XL Timothy

- **Early maturity blend**
- **Excellent with clover or alfalfa for hay or pasture**
- **1 - 2 weeks earlier to boot stage than Climax in most environments**

Top Tim XL timothy is an early maturing blend of timothy grasses. It is highly productive in northern geographies for hay or haylage. Early growth is highly palatable for pasture, while later season growth produces leafy and fine-stemmed plants with strong yields and optimum forage quality. Performs best on well drained, fertile soils.

XL The XL designation represents branded products that meet the Forage First® high-quality standard. XL brands contain one or more improved varieties.

FORAGE FIRST® FACTOR: Improved varieties of timothy are about improving its faults. Early maturing varieties align more closely when paired with alfalfa's harvest schedules. Timothy's shallow root system can struggle in warm and droughty environments. Increasing seeding rates can compensate for timothy's slow establishment, increasing stand density and weed suppression. What it lacks in vigor, it makes up in winter hardiness.

SEEDING RATES (LBS/ACRE)

MONOCULTURE
MIX COMPONENT

8 - 15 SEEDS/LB
2 - 6 DEPTH (IN)
EMERGENCE (DAYS)

1,152,000
1/4 - 1/2
14 - 21

CHARACTERISTICS

ESTABLISHMENT
PERSISTENCE
DROUGHT TOLERANCE
WINTER HARDINESS

SLOW PALATABILITY
MED YIELD POTENTIAL
LOW GRAZING TOLERANCE
HIGH

HIGH
MED
MED

PLANTING TIMES

SPRING PLANTING
FALL PLANTING

MAR - MAY LIFE CYCLE
AUG - SEP

PERENNIAL

ADAPTATION

Timothy is adapted to a cool and humid climate. Timothy thrives best on rich, moist bottomlands and on finer textured soils, such as clay loams. It does not do well on coarser soils. It prefers a pH of 5.5 to 7.0. Timothy will grow for a time on soils low in fertility, but it is better adapted to a high fertility soil. It is not well adapted to wet, flat land where water stands for any considerable time. Under limited moisture conditions, it makes a poor recovery and does not tolerate drought or prolonged high temperatures. Timothy is very winter-hardy and has high tolerance to cold temperatures and ice encasement.

ESTABLISHMENT

Timothy is usually seeded in mixes with legumes. This mix may be drilled with a small grain drill. If planted with a winter grain, the timothy is seeded with it, and the legume is planted early the following spring. A firm, weed-free seedbed is a key to a successful planting. Common seeding rates are 6 to 12 lbs/acre when seeded alone and 2 to 6 lbs/acre when seeded in mixes.

HARVEST MANAGEMENT

Cut boot to early head for first cut, then every 4 to 6 weeks thereafter.

Timothy is highly responsive to fertilizers, which should be applied frequently in ample quantities. Fertilizer, especially nitrogen, is important when legumes have almost disappeared from the hay or pasture mix. Timothy stands become weak under close and continuous grazing. A fundamental reason for the decline of timothy under poor grazing practices is injury to the corms. These corms form in the spring at the same time the stem elongates. Food materials are stored in them, and they may be destroyed by trampling of grazing animals. Timothy can be initially grazed before jointing and again between early head to full head. Second and successive grazing should also occur before jointing and when basal sprouts appear at the soil surface. After the second grazing, plants usually do not joint. Timothy should be cut for hay or silage from early to full head. Make successive harvests for hay and silage when basal sprouts appear at the soil surface. Sterile seed heads may be 15 to 20 inches up the stems when sprouts appear at the time of second cutting. Growing points stay below ground after a second cutting. Graze or cut to a minimum height of 3 inches or more.



Brought to you by
LA CROSSE
SEED
800.356.SEED
info@laxseed.com
foragefirst.com